



ORIGINAL PAPER

Correlation of the end-tidal CO₂ value with arterial blood gas parameters – evaluation of the treatment efficacy of COPD exacerbation in the emergency department

Fatma Çakmak ¹, Erdal Tekin ²

¹ Department of Emergency Medicine, Health Sciences University, Erzurum Regional Training and Research Hospital, Erzurum, Turkey

² Department of Emergency Medicine, Ataturk University Faculty of Medicine, Erzurum, Turkey

ABSTRACT

Introduction and aim. Painful, invasive, and expensive arterial blood gas (ABG) analysis is required in the diagnosis, follow-up, treatment, and even discharge of patients with chronic obstructive pulmonary disease (COPD). This study aimed to compare the end-tidal carbon dioxide (ETCO₂) value, which allows non-invasive, painless, low-cost, and continuous monitoring, with ABG parameters, in the evaluation of the treatment efficacy of COPD exacerbation.

Material and methods. The study was prospectively conducted with patients who presented to the emergency department with COPD exacerbation. ABG analysis and ETCO₂ measurement were simultaneously performed in patients with COPD exacerbation at the time of arrival and after treatment, and were statistically compared.

Results. The study included a total of 216 patients, of whom 57.4% were male. The mean age of the patients was 67.3±13.9 years. The ETCO₂ values of the patients at arrival and after COPD exacerbation treatment were 39.2±10 and 37.3±9, respectively, and a statistically significant difference was determined (p=0.001). The partial pressure of pCO₂ values measured at arrival and after treatment were 40.85±10.54 and 38.74±9.25, respectively, and it was statistically significant (p=0.001). A strong positive and statistically significant correlation was found between the ETCO₂ and pCO₂ values both at arrival and after COPD exacerbation treatment (r=0.840; p<0.001 and r=0.872; p<0.001, respectively). The Bland-Altman plot was constructed for the agreement between ETCO₂ and pCO₂ at both evaluation times.

Conclusion. ETCO₂ measurement could accurately predict the pCO₂ of patients with COPD at arrival and after COPD exacerbation treatment. Also, ETCO₂ may be useful in cases where pCO₂ cannot be used.

Keywords. arterial blood gas, capnograph, chronic obstructive pulmonary disease, end tidal carbon dioxide, partial pressure of carbon dioxide

Introduction

Chronic obstructive pulmonary disease (COPD) is a leading public health problem worldwide. Although COPD is a slowly progressive, irreversible disease, it is important because of its preventable nature and the possibility of stopping of the progression.¹⁻³ However,

COPD exacerbations adversely affect the patient's health status, hospitalization, and length of stay.⁴ In these patients, arterial blood gas (ABG) measurement is one of the valuable tests used to determine the patient's oxygenation, ventilation, and acid-base status during and after an exacerbation. However, ABG is an invasive and

Corresponding author: Erdal Tekin, e-mail: dret25@gmail.com

Received: 23.05.2022 / Revised: 19.06.2022 / Accepted: 21.06.2022 / Published: 30.09.2022

Çakmak F, Tekin E. Correlation of the end-tidal CO₂ value with arterial blood gas parameters – evaluation of the treatment efficacy of COPD exacerbation in the emergency department. *Eur J Clin Exp Med*. 2022;20(3):316–322. doi: 10.15584/ejcem.2022.3.9.



painful procedure and may need to be repeated in certain cases. Therefore, alternative methods have been used in the evaluation of diagnosis, and follow-up of these patients.^{5,6} One of these alternative methods is the measurement of end-tidal carbon dioxide (ETCO₂).

ETCO₂ is determined by non-invasively with a capnometer by measuring the partial pressure of expiratory carbon dioxide from the air inhaled and exhaled during respiration.⁷ This parameter is generally used in the emergency department (ED) to confirm the location of the endotracheal tube, monitor spontaneous return during cardiopulmonary resuscitation in patients with cardiac arrest, and monitor ventilation status in procedural sedoanalgesia.^{8,9} In addition, it can provide instant information concerning the metabolic status, perfusion, and most importantly ventilation of critically ill patients in the ED.⁴ Studies have emphasized that ETCO₂ can be used to evaluate the severity of obstructive airway disease in the ED.^{7,10}

Aim

In this study, we aimed to compare the ETCO₂ value measured simultaneously with the ABG in patients who came to the ED with COPD exacerbation, at the time of admission and after COPD exacerbation treatment was applied.

Material and methods

Study design

This single-center, prospective clinical study was conducted with patients who presented to the ED with COPD exacerbation, between June 27th, 2019 and December 27th, 2019. The tertiary university hospital, where the study was carried out, provides health services to approximately 4.5 million people in 12 neighboring provinces. The study protocol was in accordance with the Declaration of Helsinki and Good Clinical Practices. Informed consent was obtained from all patients participating in the study. Local ethics committee approval was obtained for the study (decision number: 27.06.2019/05-28).

Patient selection

The eligible participants and sample population were the patients who were admitted to the ED with COPD exacerbations during the study period. Patients with shortness of breath due to reasons other than COPD exacerbation, those younger than 18 years, those with contraindications to ABG sampling (such as coagulopathy, local infection, and thrombus), those that did not provide consent for participation in the study, pregnant patients, intubated patients or patients that required intubation during their follow-up, patients who could not tolerate capnometry for the ETCO₂ measurement, those with anatomical disabilities, and non-compliant patients were excluded from the study.

ABG sampling

In the ABG sampling procedure, after the radial artery pulse of the patient was palpated, the most suitable site for the intervention was determined and cleaned with 10% povidone iodine. Then, arterial puncture was performed by the blood gas determination set containing lithium heparin. All interventions in the ED were undertaken by the same physician. The ADL 800 Flex device was used for the ABG analyses. The blood gas analyzer was calibrated according to the manufacturer's recommendations.

Study protocol

Patients presenting to the ED and included in the study were asked to breathe through a nasal cannula connected to a capnometer for at least 30 seconds, during which ABG samples were obtained. The ETCO₂ value was simultaneously recorded with ABG parameters. The ETCO₂ measurement was performed non-invasively by the same physician using a capnograph/pulse oximeter device, Capnostream®20p portable bedside monitor. To make an accurate measurement, the patient was monitored for at least 6 minutes (240 seconds) with a capnograph/pulse oximeter.¹¹ During the procedure, SpO₂, rhythm (rate and rhythm) and ETCO₂ were monitored and respiratory rate were recorded. Then, COPD exacerbation treatment was applied to the patients in line with the guidelines of the Global Initiative on Chronic Obstructive Pulmonary Disease.¹² After the initial evaluation and exacerbation treatment, the patients were asked to breathe again with a capnometer, and the ETCO₂ value, respiratory rate, body temperature, heart rate, arterial blood pressure, and oxygen saturation values were recorded again simultaneously with the ABG parameters.

Treatment efficacy

The efficacy of treatment was evaluated by comparing vital signs, ABG parameters, and ETCO₂ measured at the time of arrival at the ED and after COPD exacerbation treatment.

Outcome measures

The primary outcome of the study was the correlation between ABG parameters and ETCO₂, which were obtained at the time of arrival at the ED and after COPD exacerbation treatment. The secondary outcome was the treatment efficacy evaluated by comparing the pre-treatment and post-treatment vital signs, ABG parameters, and ETCO₂.

Statistical analysis

All statistical analyses were performed using the Statistical Package for the Social Science (SPSS), Version 20.0 (IBM Corp., Armonk, New York, USA). Data were

presented as mean, standard deviation, percentage and frequency. The normality of the distribution of continuous variables was analyzed by using the Shapiro-Wilk test. In the comparisons of two dependent groups, the paired-samples t-test was used for the normally distributed data, and the Wilcoxon test for data without a normal distribution. In the comparison of two continuous variables, the Pearson correlation test was conducted if the normal distribution was determined, and the Spearman correlation test otherwise. The agreement between the ETCO_2 value and ABG parameters was evaluated by using the Bland-Altman method. The statistical significance level was taken as $p<0.05$.

Results

Patient population and characteristics

During the study, a total of 556 patients were presented to the ED due to COPD exacerbation. After applying the inclusion and exclusion criteria, a total of 216 patients were included in the study. The flow chart of the study is given in Figure 1.




Fig. 1. Flow diagram of the study

Table 1 presents the demographic and clinical features of the patients. The mean age of the patients was 67.3 ± 13.9 years, and 124 (57.4%) were male. All of the patients included in the study were COPD patients. The second most common disease detected in the participants was hypertension ($n=107$, 49.5%).

Comparison of pre-treatment and post-treatment evaluations

Figure 2 presents the comparison of the clinical features, ETCO_2 values and ABG parameters of the patients at the time of arrival and after COPD exacerbation treatment. There was a statistical significance between the pre- and post-treatment vital signs, ETCO_2 values and ABG parameters ($p<0.05$).

Table 1. Demographic and clinical characteristics of the patients ($n=216$)^a

Variable	Value
Mean age (years)	67.3 ± 13.9
Gender	
Male, n (%)	124 (57.4%)
Female, n (%)	92 (42.6%)
Condenser use, n (%)	110 (50.9%)
Smokers, n (%)	191 (88.4%)
Smoking, pack-day	1.2 ± 0.1
Clinical characteristics	
Respiratory rate (breaths/min)	25.1 ± 7.2
Body temperature (°C)	36.9 ± 0.3
Heart rate (beats/min)	99.8 ± 19.9
Systolic arterial blood pressure (mmHg)	139.9 ± 19.9
Diastolic arterial blood pressure (mmHg)	84.2 ± 12.3
Oxygen saturation (%)	80.1 ± 10.6
ETCO_2	39.2 ± 10
Chronic diseases other than COPD	
Hypertension, n (%)	107 (49.5%)
Diabetes mellitus, n (%)	69 (31.9%)
Congestive heart failure, n (%)	56 (25.9%)
Other, n (%)	43 (19.9%)

^a COPD – chronic obstructive pulmonary disease, ETCO_2 – end-tidal carbon dioxide




Fig. 2. Comparison of the patients' vital signs, ETCO_2 values and ABG parameters measured at arrival and after treatment (X: $p=0.001$, +: $p=0.002$, ++: $p=0.042$, *: $p=0.015$)

The relationship between the ETCO_2 value and ABG parameters measured at arrival is shown in Figure 3. There was a positive, strong and statistically significant correlation between ETCO_2 and pCO_2 ($r=0.840$, $p<0.001$).

Figure 4 shows the relationship between the ETCO₂ values and ABG parameters measured after COPD exacerbation treatment. Likewise, a positive, strong and statistically significant correlation was found between ETCO₂ and pCO₂ ($r=0.872$, $p<0.001$). According to the Bland-Altman analysis, the mean \pm standard deviation of the difference between the pCO₂ and ETCO₂ values measured at the time of arrival was 1.64 ± 5.73 [95% confidence interval (CI): 0.87-2.41], and the upper and lower limits were determined as 12.87 and -9.59, respectively (Figure 5a). In the Bland-Altman analysis of the difference between the post-treatment pCO₂ and ETCO₂, the mean \pm standard deviation was found to be 1.45 ± 4.41 (95% CI: 0.86-2.04), and the upper and lower limits were 10.09 and -7.19, respectively (Figure 5b).

Discussion

In this study, the ETCO₂ value was compared with ABG parameters in patients applied with COPD exacerbation, at the time of arrival and after the treatment. In order to determine the efficacy of COPD exacerbation treatment, the vital signs and ABG parameters of the patients were examined before and after treatment, and the differences were found to be statistically significant. When the correlation of the ETCO₂ value with the ABG parameters was evaluated, ETCO₂ had a strong correlation with pCO₂ both at arrival and after treatment, and had a moderately positive and statistically significant correlation with HCO₃. In addition, there was a clear agreement between ETCO₂ and pCO₂ both before and after treatment according to the Bland-Altman graph.

ABG evaluation is of great importance in the hospitalization and discharge decisions after the treatment of patients with COPD exacerbation. The ABG analysis is an invasive and painful procedure that may be accompanied by possible complications.¹³ Therefore, the analysis of ABG is difficult in some cases.¹⁴ During the study, it was also observed that patients did not want to give an ABG sample for the second time.

Capnography, which is a non-invasive, painless and uncomplicated method, provides instant information concerning the patient's status by measuring the ETCO₂ value and respiratory rate in patients with acute respiratory problems, such as COPD exacerbations.¹⁵ In the literature, comparisons have been made between ETCO₂ and pCO₂ in the evaluation of respiratory status of the patients who have undergone sedation and general anesthesia, except for COPD cases, and the results have been reported to be statistically significantly.^{8,16-18} In the current study, it was shown that the ETCO₂ value could be used in the follow-up of patients with COPD and in the evaluation of the efficacy of COPD exacerbation treatment. Dogan et al., who investigated the accuracy of the ETCO₂ level in the prediction of the severity of COPD exacerbation in the ED, found the presence of a relationship between ETCO₂ and pCO₂, and pO₂, similar to the results of our study.¹⁰

Cinar et al. investigated whether there was a difference between the ABG pCO₂ value and the ETCO₂ value in patients applied to the ED due to acute dyspnea, and reported a positive, strong and statistically significant correlation between these two parameters ($r=0.911$; $p<0.001$).¹⁹ The authors also emphasized that




Fig. 3. Correlation of the patients' ETCO₂ values with arterial blood gas parameters at arrival




Fig. 4. Correlation of the patients' post-treatment ETCO₂ values with arterial blood gas parameters




Fig. 5. Bland-Altman graph (a. agreement between ETCO₂ and pCO₂ at arrival, b. agreement between ETCO₂ and pCO₂ after COPD exacerbation treatment)

ETCO₂ could accurately predict the pCO₂ value of patients presenting to the ED with acute dyspnea. In another study conducted with adult asthmatic patients with acute exacerbation, Corbo et al. found a significant correlation between pCO₂ in blood gas and ETCO₂ measured by capnography.²⁰ Kartal et al. investigated the value of ETCO₂ measurement for patients with COPD in the ED and concluded that there was a moderate correlation between the ETCO₂ and pCO₂ levels, but unlike other studies, ETCO₂ measurement was not sufficient to predict the pCO₂ level in this patient group.⁴ Similarly, Pekdemir et al. reported that the ETCO₂ value was significantly lower than the pCO₂ value and that there was a weak correlation between these two parameters.²¹ In a

review of ETCO₂, it was emphasized that, ETCO₂ was an important tool in evaluating the severity of obstructive airway disease.⁷ In acute respiratory problems, ETCO₂ can detect airway obstruction and hypoventilation at an early stage. Thus, they are able to provide earlier medical intervention to these patients.

Taghizadieh et al. compared ETCO₂ with bicarbonate (HCO₃), an ABG parameter, and emphasized that, although ETCO₂ could be used in the primary diagnosis of metabolic acidosis, ABG was still the gold standard for guiding diagnosis and treatment.²² However, Kartal et al. suggested that the ETCO₂ level could indicate the severity of metabolic acidosis and mortality.⁴ In another study comparing ETCO₂ with HCO₃ in pa-

tients with COPD exacerbation, Taghizadieh et al. detected no statistically significant difference between the ET-CO₂ and HCO₃ levels ($r=0.04$; $p=0.136$).²³ In the same study, the authors found a low negative correlation between ET-CO₂ and pH ($r=-0.249$; $p<0.001$), and a moderate positive correlation between ET-CO₂ and pCO₂ ($r=0.611$; $p<0.001$). In the current study, the ET-CO₂ and HCO₃ levels of the patients were positively correlated both at arrival and after treatment, and this correlation was statistically significant. In addition, we observed a low level of negative correlation between ET-CO₂ and pH both at arrival and after treatment. According to the data obtained from our study, there was a strong positive correlation, similar distribution and agreement between the values of ET-CO₂ and PCO₂ measured at the time of arrival. More importantly, this correlation was even stronger in the post-treatment ET-CO₂ and PCO₂ measurements of the patients, showing less bias and lower difference, and a much better agreement.

Limitations

The first limitation of the study was that it was conducted in a single center. Secondly, patients with concomitant metabolic diseases, including diabetes mellitus and acute renal failure, which affect metabolic parameters, such as pH and HCO₃, were not excluded from the sample. Since real-world COPD patients may have some other comorbidities, and our study aimed to study all COPD patients, patients with or without certain concomitant metabolic diseases, such as diabetes or acute renal failure, were not excluded. Subsequently, factors such as cardiomegaly and elevated pulmonary artery pressure that could affect the correlation between pCO₂ and ET-CO₂ were not evaluated. Also, body temperature, which may affect the pCO₂ level, was not taken into consideration. Lastly, the cigarette consumption of the patients was calculated as package per day. This, in turn, does not report how much COPD patients actually consume package cigarettes per year.

Conclusion

It was determined that ET-CO₂ measurement which was non-invasive, painless, repeatable and low-cost capnography procedure may provide information about the ventilation, circulation and metabolism of patients during COPD exacerbation at the time of arrival and after treatment. Considering the similar and strongly related properties of ET-CO₂ to PCO₂ and HCO₃, we believe that it may be used during the follow-up of patients.

Declarations

Funding

The authors declared that this study has received no financial support or any funding.

Author contributions

Conceptualization, F.Ç. and E.T.; Methodology, F.Ç. and E.T.; Software, F.Ç.; Validation, F.Ç. and E.T.; Formal Analysis, F.Ç. and E.T.; Investigation, F.Ç.; Resources, F.Ç.; Data Curation, F.Ç. and E.T.; Writing – Original Draft Preparation, F.Ç. and E.T.; Writing – Review & Editing, F.Ç. and E.T.; Visualization, F.Ç.; Supervision, E.T.

Conflicts of interest

The authors have no conflict of interest to declare.

Data availability

The data that support the findings of this study are available from the corresponding author, upon reasonable request.

Ethics approval

Local ethics committee approval was obtained from the Ethics Committee of Faculty of Medicine, University of Ataturk, Erzurum, Turkey, with the decision number of 27.06.2019/05-28.

References

- Stanford RH, Korner S, Brekke L, Reinsch T, Bengtson LG. Validation and assessment of the COPD treatment ratio as a predictor of severe Exacerbations. *Chronic Obstr Pulm Dis.* 2020;7(1):38-48.
- Vogelmeier CF, Román-Rodríguez M, Singh D, Han MK, Rodríguez-Roisin R, Ferguson GT. Goals of COPD treatment: Focus on symptoms and exacerbations. *Respir Med.* 2020;166:105938.
- Arslan S, Delice O, Katran ZY. The predictive value of qSOFA and CURB-65 scores in predicting the hospitalization need and mortality in patients with acute exacerbation of chronic obstructive pulmonary disease. *Ital J Emerg Med.* 2020;9(3):169-177.
- Kartal M, Goksu E, Eray O, et al. The value of ET-CO₂ measurement for COPD patients in the emergency department. *Eur J Emerg Med.* 2011;18(1):9-12.
- Mummery V, Rogers E, Padmanaban V, Matthew D, Woodcock T, Bloch S. Transcutaneous carbon dioxide measurement is not a reliable alternative to arterial blood gas sampling in the acute medical setting. *Eur Respir J.* 2019;53(4):1801726.
- Huttmann SE, Windisch W, Storre JH. Techniques for the measurement and monitoring of carbon dioxide in the blood. *Ann Am Thorac Soc.* 2014;11(4):645-652.
- Aminiahidashti H, Shafiee S, Zamani Kiasari A, Sazgar M. Applications of End-Tidal Carbon Dioxide (ET-CO₂) Monitoring in Emergency Department; a Narrative Review. *Emerg (Tehran).* 2018;6(1):e5.
- Long B, Koyfman A, Vivirito MA. Capnography in the Emergency Department: A Review of Uses, Waveforms, and Limitations. *J Emerg Med.* 2017;53(6):829-842.

9. Cereceda-Sánchez FJ, Molina-Mula J. Capnography as instrument for airway management during basic instrumental CPR manoeuvres. *Resuscitation*. 2019;138:132–133.
10. Dogan NO, Sener A, Gunaydin GP, et al. The accuracy of mainstream end-tidal carbon dioxide levels to predict the severity of chronic obstructive pulmonary disease exacerbations presented to the ED. *Am J Emerg Med*. 2014;32(5):408–411.
11. Galia F, Brimiouille S, Bonnier F, et al. Use of Maximum End-Tidal CO₂ Values to Improve End-Tidal CO₂ Monitoring Accuracy. *Respir Care*. 2011;56(3):278–283.
12. Global Strategy for Diagnosis, Management and Prevention of COPD. The Global Initiative for Chronic Obstructive Lung Diseases (GOLD). 2020 report. <https://goldcopd.org/gold-reports/>. Accessed September 24, 2021.
13. Tyagi D, Govindagoudar MB, Jakka S, Chandra S, Chaudhry D. Correlation of PaCO₂ and ETCO₂ in COPD Patients with Exacerbation on Mechanical Ventilation. *Indian J Crit Care Med*. 2021;25(3):305–309.
14. Ekström M, Engblom A, Ilıc A, Holthius N, Nordström P, Vaara I. Calculated arterial blood gas values from a venous sample and pulse oximetry: Clinical validation. *PLoS One*. 2019;14(4):e0215413.
15. Schwarz SB, Windisch W, Magnet FS, et al. Continuous non-invasive PCO₂ monitoring in weaning patients: Transcutaneous is advantageous over end-tidal PCO₂. *Respirology*. 2017;22(8):1579–1584.
16. Bhavani-Shankar K, Philip JH. Defining segments and phases of a time capnogram. *Anesth Analg*. 2000;91(4):973–977.
17. Onodi C, Bühler PK, Thomas J, Schmitz A, Weiss M. Arterial to end-tidal carbon dioxide difference in children undergoing mechanical ventilation of the lungs during general anaesthesia. *Anaesthesia*. 2017;72(11):1357–1364.
18. Kummer DJ, Walden BJ. Capnography Monitoring for Patients Undergoing Moderate Sedation: An SGNA Fellowship Project. *Gastroenterology Nursing*. 2019;42(1):49–54.
19. Cinar O, Acar YA, Arziman İ, Kilic E, Eyi YE, Ocal R. Can mainstream end-tidal carbon dioxide measurement accurately predict the arterial carbon dioxide level of patients with acute dyspnea in ED. *Am J Emerg Med*. 2012;30(2):358–361.
20. Corbo J, Bijur P, Lahn M, Gallagher EJ. Concordance between capnography and arterial blood gas measurements of carbon dioxide in acute asthma. *Ann Emerg Med*. 2005;46(4):323–327.
21. Pekdemir M, Cinar O, Yilmaz S, Yaka E, Yuksel M. Disparity between mainstream and sidestream end-tidal carbon dioxide values and arterial carbon dioxide levels. *Respir Care*. 2013;58(7):1152–1156.
22. Taghizadieh A, Pouraghaei M, Moharamzadeh P, Ala A, Rahmani F, Basiri Sofiani K. Comparison of end-tidal carbon dioxide and arterial blood bicarbonate levels in patients with metabolic acidosis referred to emergency medicine. *J Cardiovasc Thorac Res*. 2016;8(3):98–101.
23. Taghizadieh A, Rahmani F, Soleimanpour H, Aminifazl H. Comparison of end tidal carbon dioxide and arterial blood bicarbonate levels in patients with exacerbation chronic obstructive pulmonary disease. *Thrita*. 2015;4(2):e26169.